

PAPER-1: MATHEMATICS, CHEMISTRY & PHYSICS**A5****Do not open this test booklet until you are asked to do so****Read carefully the Instruction on the Back Cover of this Test Booklet.****IMPORTANT INSTRUCTIONS**

1. Immediately fill in the particulars on this page of the Test Booklet with Blue/Black Point Pen. *Use of pencil is strictly prohibited.*
- 2.. The answer sheet is kept inside this test Booklet. When you are directed to open the Test Booklet, take out the Answer sheet and fill in the particulars carefully.
3. The test is of **3 hours** duration.
4. The Test Booklet consists of **105 questions of 3 marks each**. The maximum marks are **315**.
5. There are three parts in the question paper.
The distribution of marks subjectwise in each part is as under for each correct response.
Part-A - Mathematics (105 marks) - 35 questions.
Part-B - Chemistry (105 marks) - 35 questions.
Part-C - Physics (105 marks) - 35 questions.
6. *Candidates will be awarded three marks each for indicated correct response of each question. **One mark** will be deducted for indicated incorrect response of each question. **No deduction** from the total score will be made **if no response** is indicated for an item in the Answer Sheet.*
7. Use **Blue/Black Ball point pen only** for writing particulars/markings responses on **Side-1** and **Side-2** of the answer sheet. **Use of pencil is strictly prohibited.**
8. No candidate is allowed to carry any textual material, printed or written, bits of papers, pager, mobile phone, any electronic device, etc., except the Admit card inside the examination hall/room.
9. Rough work is to be done on the space provided for this purpose in the Test Booklet only. This space is given at the bottom of each page and in 4 pages at the end of the booklet.
10. On completion of the test, the candidate must hand over the Answer Sheet to the Invigilator on duty in the Room/Hall. **However, the candidates are allowed to take away this Test Booklet with them.**
11. The CODE for this booklet is **A5**. Make sure that the CODE printed on **Side-2** of the answer sheet is the same as that on this booklet. In case of discrepancy, the candidate should immediately report the matter to the Invigilator for replacement of both the Test Booklet and the Answer Sheet.
12. **Do not fold or make any stray marks on the Answer Sheet.**

Name of the Candidate (in Capital Letters): _____

Roll Number : in figure _____

: in words _____

Examination Centre Number: _____

Name of Examination Centre (in Capital Letters) : _____

Candidate's Signature : _____ Invigilator Signature _____

PART-A: MATHEMATICS

1. Let $f : N \rightarrow Y$ be a function defined as $f(x) = 4x + 3$ where $Y = \{y \in N : y = 4x + 3 \text{ for some } x \in N\}$. Show that f is invertible and its inverse is

$$(1) \quad g(y) = 4 + \frac{y+3}{4} \quad (2) \quad g(y) = \frac{y+3}{4} \quad (3) \quad g(y) = \frac{y-3}{4} \quad (4) \quad g(y) = \frac{3y+4}{3}$$

Sol: Ans [3]

$f(x)$ is both one-one and onto

\Rightarrow It is invertible

Now, let $f(x) = y = 4x + 3$

$$\Rightarrow x = \frac{y-3}{4}$$

$$f^{-1}(y) = \frac{y-3}{4}$$

$$\Rightarrow g(y) = \frac{y-3}{4}$$

2. Let R be the real line. Consider the following subsets of the plane $R \times R$:

$$S = \{(x, y) : y = x + 1 \text{ and } 0 < x < 2\}$$

$$T = \{(x, y) : x - y \text{ is an integer}\}.$$

Which one of the following is true?

- (1) Both S and T are equivalence relations on R
- (2) S is an equivalence relation on R but T is not
- (3) T is an equivalence relation on R but S is not
- (4) Neither S nor T is an equivalence relation on R

Sol: Ans [3]

Reflexive: as $x \neq x + 1 \Rightarrow S$ is not reflexive $\Rightarrow S$ is not equivalence

as $x - x$ is an integer $\Rightarrow T$ is reflexive

Symmetric:

$(x - y)$ is an integer $\Rightarrow (y - x)$ is an integer

$\Rightarrow T$ is symmetric

Transitive:

Let $x - y$ be an integer, $y - z$ be an integer

$$x - y = n_1 \quad y - z = n_2$$

$$\Rightarrow x - y + y - z = n_1 + n_2$$

$$\Rightarrow (x - z) \text{ is an integer}$$

$$\Rightarrow T \text{ is transitive} \Rightarrow T \text{ is equivalence.}$$

3. The conjugate of complex number is $\frac{1}{i-1}$. Then that complex number is

(1) $\frac{1}{i+1}$

(2) $\frac{-1}{i+1}$

(3) $\frac{1}{i-1}$

(4) $\frac{-1}{i-1}$

Sol: Ans [2]

$$\text{Let } z = \frac{1}{i-1}$$

$$\therefore \bar{z} = \frac{1}{-i-1} = \frac{-1}{i+1}$$

4. The quadratic equations

$$x^2 - 6x + a = 0$$

$$\text{and } x^2 - cx + 6 = 0$$

have one root in common. The other roots of the first and second equations are integers in the ratio 4 : 3. Then the common root is

(1) 4

(2) 3

(3) 2

(4) 1

Sol: Ans [3]

$$x^2 - 6x + a = 0 \quad \text{let roots be } \alpha, \beta$$

$$x^2 - 6x + 6 = 0 \quad \text{let roots be } \alpha, \gamma$$

$$\alpha\beta = a \quad \alpha\gamma = 6$$

$$\Rightarrow \frac{\beta}{\gamma} = \frac{a}{6} = \frac{4}{3} \quad \Rightarrow a = 8$$

$$\Rightarrow \alpha^2 - 6\alpha + 8 = 0$$

$$\Rightarrow \alpha^2 - 4\alpha - 2\alpha + 8 = 0$$

$$\Rightarrow \alpha(\alpha - 4) - 2(\alpha - 4) = 0$$

$$\Rightarrow \alpha = 2, 4$$

$$\text{if } \alpha = 4,$$

$$\text{then } \alpha \cdot \gamma = 6$$

$$\Rightarrow \gamma \text{ is not an integer}$$

$$\Rightarrow \alpha = 2$$

5. Let A be a square matrix all of whose entries are integers. Then which one of the following is true?

(1) If $\det A \neq \pm 1$, then A^{-1} exists and all its entries are non-integers

(2) If $\det A = \pm 1$, then A^{-1} exists and all its entries are integers

(3) If $\det A = \pm 1$, then A^{-1} need not exist

(4) If $\det A = \pm 1$, then A^{-1} exists but all its entries are not necessarily integers

Sol: Ans [2]

If $\det A \neq 0 \Rightarrow A^{-1}$ exists

and $A^{-1} = \frac{\text{adj } A}{|A|}$

$\therefore \text{adj } A = \text{Transpose of cofactor matrix of } A$. If A has integral entries, then $\text{adj } A$ will also have integral entries and if $\det(A) = \pm 1$, then A^{-1} will also have integral entries.

6. Let a, b, c be any real numbers. Suppose that there are real numbers x, y, z not all zero such that $x = cy + bz, y = az + cx$ and $z = bx + ay$. Then $a^2 + b^2 + c^2 + 2abc$ is equal to

- (1) -1 (2) 0 (3) 1 (4) 2

Sol: Ans [3]

$$x - cy - bz = 0$$

$$cx - y + az = 0$$

$$bx + ay - z = 0$$

for non zero solutions, $|A| = 0$

$$\begin{vmatrix} 1 & -c & -b \\ c & -1 & a \\ b & a & -1 \end{vmatrix} = 0$$

$$\Rightarrow 1(1 - a^2) + c(-c - ab) - b(ac + b) = 0$$

$$\Rightarrow 1 - a^2 - c^2 - abc - b^2 - abc = 0$$

$$\Rightarrow a^2 + b^2 + c^2 + 2abc = 1$$

7. How many different words can be formed by jumbling the letters in the word MISSISSIPPI in which no two S are adjacent?

- (1) $6 \cdot 7 \cdot {}^8C_4$ (2) $6 \cdot 8 \cdot {}^7C_4$ (3) $7 \cdot {}^6C_4 \cdot {}^8C_4$ (4) $8 \cdot {}^6C_4 \cdot {}^7C_4$

Sol: Ans [3]

The number of ways $= {}^8C_4 \times \frac{7!}{4! 2!} = {}^8C_4 \cdot \frac{7 \times 6 \times 5}{2 \times 1} = {}^8C_4 \cdot 7 \cdot {}^6C_4$

8. The first two terms of a geometric progression add upto 12. The sum of the third and the fourth terms is 48. If the terms of the geometric progression are alternately positive and negative, then the first term is

- (1) -12 (2) 12 (3) 4 (4) -4

Sol: Ans [1]

$$a + ar = 12 \quad ar^2 + ar^3 = 48$$

$$a(1 + r) = 12 \quad ar^2(1 + r) = 48$$

$$\Rightarrow \frac{ar^2(1 + r)}{a(1 + r)} = \frac{48}{12}$$

$$\Rightarrow r = -2 \quad (\text{as terms are alternately positive and negative})$$

$$\Rightarrow a(1 - 2) = 12$$

$$\Rightarrow a = -12$$

9. Let $f(x) = \begin{cases} (x-1) \sin \frac{1}{x-1} & \text{if } x \neq 1 \\ 0 & \text{if } x = 1 \end{cases}$. Then which one of the following is true?

- (1) f is differentiable at $x = 0$ and at $x = 1$
- (2) f is differentiable at $x = 0$ but not at $x = 1$
- (3) f is differentiable at $x = 1$ but not at $x = 0$
- (4) f is neither differentiable at $x = 0$ nor at $x = 1$

Sol: Ans [2]

Using algebra of differentiable functions,

At $x = 0$,

$(x - 1)$ is differentiable and $\sin\left(\frac{1}{x-1}\right)$ is differentiable. So $(x - 1) \cdot \sin\left(\frac{1}{x-1}\right)$ is differentiable

At $x = 1$,

$$f'(1^-) = \lim_{h \rightarrow 0^+} \frac{(1-h-1) \sin\left(\frac{1}{1-h-1}\right) - 0}{-h} = \lim_{h \rightarrow 0^+} \sin\left(-\frac{1}{h}\right)$$

= any number between -1 to 1

⇒ LHD is oscillating

⇒ $f(x)$ is non differentiable at $x = 1$.

10. How many real solutions does the equation $x^7 + 14x^5 + 16x^3 + 30x - 560 = 0$ have?

- (1) 1
- (2) 3
- (3) 5
- (4) 7

Sol: Ans [1]

$$f(x) = x^7 + 14x^5 + 16x^3 + 30x - 560$$

+ + + + -

Change of sign ⇒ max 1 positive real root

$$f(-x) = -x^7 - 14x^5 - 16x^3 - 30x - 560$$

⇒ no change of sign ⇒ no negative real root

⇒ only 1 real solution.

11. Suppose the cubic $x^3 - px + q$ has three distinct real roots where $p > 0$ and $q > 0$. Then which one of the following holds?

- (1) The cubic has minima at $-\sqrt{\frac{p}{3}}$ and maxima at $\sqrt{\frac{p}{3}}$
- (2) The cubic has minima at both $\sqrt{\frac{p}{3}}$ and $-\sqrt{\frac{p}{3}}$
- (3) The cubic has maxima at both $\sqrt{\frac{p}{3}}$ and $-\sqrt{\frac{p}{3}}$
- (4) The cubic has minima at $\sqrt{\frac{p}{3}}$ and maxima at $-\sqrt{\frac{p}{3}}$

Sol: Ans [4]

Let $f(x) = x^3 - px + q$

$f'(x) = 3x^2 - p = 0$

$\Rightarrow x = \pm \sqrt{\frac{p}{3}}$

Now $f''(x) = 6x$

$f''\left(\sqrt{\frac{p}{3}}\right) > 0$ so minima at $x = \sqrt{\frac{p}{3}}$

$f''\left(-\sqrt{\frac{p}{3}}\right) < 0$ so maxima at $x = -\sqrt{\frac{p}{3}}$

12. The value of $\sqrt{2} \int \frac{\sin x}{\sin(x - \pi/4)} dx$ is

(1) $x - \log |\sin(x - \pi/4)| + c$

(2) $x + \log |\sin(x - \pi/4)| + c$

(3) $x - \log |\cos(x - \pi/4)| + c$

(4) $x + \log |\cos(x - \pi/4)| + c$

Sol: Ans [2]

$\sqrt{2} \int \frac{\sin x}{\sin(x - \pi/4)} dx$

Let $x - \frac{\pi}{4} = t \Rightarrow dx = dt$

$\Rightarrow \int \left(\frac{\sin t + \cos t}{\sin t} \right) dt$

$\Rightarrow \int (1 + \cot t) dt$

$\Rightarrow t + \log |\sin t| + c$

$\Rightarrow x + \log |\sin(x - \pi/4)| + c$

13. The area of the plane region bounded by the curves $x + 2y^2 = 0$ and $x + 3y^2 = 1$ is equal to

(1) $\frac{1}{3}$

(2) $\frac{2}{3}$

(3) $\frac{4}{3}$

(4) $\frac{5}{3}$


Sol: Ans [3]

$y^2 = -x/2$

$y^2 = -\frac{(x-1)}{3}$

$\therefore A = 2 \int_0^1 ((1 - 3y^2) - (-2y^2)) dy$

$= 2 \int_0^1 (1 - y^2) dy = \frac{4}{3}$



14. Let $I = \int_0^1 \frac{\sin x}{\sqrt{x}} dx$ and $J = \int_0^1 \frac{\cos x}{\sqrt{x}} dx$. Then which one of the following is true?

- (1) $I < \frac{2}{3}$ and $J < 2$ (2) $I < \frac{2}{3}$ and $J > 2$ (3) $I > \frac{2}{3}$ and $J < 2$ (4) $I > \frac{2}{3}$ and $J > 2$

Sol: Ans [1]

$$I = \int_0^1 \frac{\sin x}{\sqrt{x}} dx = \int_0^1 \frac{x - \frac{x^3}{3!} + \frac{x^5}{5!} - \dots}{\sqrt{x}} dx$$

$$\Rightarrow I < \int_0^1 \sqrt{x} dx = \frac{2}{3}$$

Again $J = \int_0^1 \frac{\cos x}{\sqrt{x}} dx = \int_0^1 \frac{1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \dots}{\sqrt{x}} dx$

$$J < \int_0^1 x^{-1/2} dx$$

$$J < 2$$

15. The differential equation of the family of circles with fixed radius 5 units and centre on the line $y = 2$ is

- (1) $(y - 2)y'^2 = 25 - (y - 2)^2$ (2) $(y - 2)^2y'^2 = 25 - (y - 2)^2$
 (3) $(x - 2)^2y'^2 = 25 - (y - 2)^2$ (4) $(x - 2)y'^2 = 25 - (y - 2)^2$

Sol: Ans [2]

Let centre be $(\alpha, 2)$

$$\therefore \text{Equation of circle is } (x - \alpha)^2 + (y - 2)^2 = 25 \dots(i)$$

On differentiating,

$$2(x - \alpha) + 2(y - 2)y' = 0$$

$$\therefore (x - \alpha) = -(y - 2)y'$$

Putting in (i),

$$(y - 2)^2y'^2 = 25 - (y - 2)^2$$

16. The solution of the differential equation

$$\frac{dy}{dx} = \frac{x + y}{x}$$

satisfying the condition $y(1) = 1$ is

- (1) $y = x \ln x + x^2$ (2) $y = xe^{(x-1)}$ (3) $y = x \ln x + x$ (4) $y = \ln x + x$

Sol: Ans [3]

$$\frac{dy}{dx} = 1 + \frac{y}{x}$$

$$\frac{dy}{dx} + \left(\frac{-1}{x}\right)y = 1$$

$$\text{I.F.} = e^{\int \frac{-1}{x} dx} = e^{-\ln x} = \frac{1}{x}$$

$$\Rightarrow y \cdot \frac{1}{x} = \int \left(1 \times \frac{1}{x}\right) dx + c$$

$$\Rightarrow y \cdot \frac{1}{x} = \ln x + c$$

$$\Rightarrow y = x \ln x + cx$$

$$\Rightarrow c = 1 \quad (\text{as } y(1) = 1)$$

$$\Rightarrow y = x \ln x + x$$

17. The perpendicular bisector of the line segment joining $P(1, 4)$ and $Q(k, 3)$ has y-intercept -4 . Then a possible value of k is

- (1) 2 (2) -2 (3) -4 (4) 1

Sol: Ans [3]

$$\text{Slope of } PQ = \frac{3-4}{k-1} = \frac{-1}{k-1}$$

$$\therefore \text{Slope of } RS = \frac{15}{k+1}$$


As PQ and RS are perpendicular,

$$\left(\frac{-1}{k-1}\right) \times \frac{15}{k+1} = -1$$

$$\Rightarrow k^2 - 1 = 15$$

$$\Rightarrow k^2 = 16$$

$$\Rightarrow k = 4, -4$$



18. The point diametrically opposite to the point $P(1, 0)$ on the circle $x^2 + y^2 + 2x + 4y - 3 = 0$ is

- (1) (-3, 4) (2) (-3, -4) (3) (3, 4) (4) (3, -4)

Sol: Ans [2]

$$x^2 + y^2 + 2x + 4y - 3 = 0$$

Centre $\equiv (-1, -2)$

Let point be (h, k)


$$\Rightarrow \frac{h+1}{2} = -1 \quad \frac{k+0}{2} = -2$$

$$\Rightarrow (h, k) \equiv (-3, -4)$$

19. A parabola has the origin as its focus and the line $x = 2$ as the directrix. Then the vertex of the parabola is at

- (1) (1, 0) (2) (0, 1) (3) (2, 0) (4) (0, 2)

Sol: Ans [1]



Vertex is mid-point of S and Z

⇒ vertex = (1, 0)

20. A focus of an ellipse is at the origin. The directrix is the line $x = 4$ and the eccentricity is $1/2$. Then the length of the semi-major axis is

- (1) $2/3$ (2) $4/3$ (3) $5/3$ (4) $8/3$

Sol: Ans [4]

Focus = (0, 0) $e = \frac{1}{2}$

Directrix ⇒ $x - 4 = 0$

⇒ equation of ellipse

⇒ $x^2 + y^2 = \frac{1}{4}(x - 4)^2$

⇒ $4x^2 + 4y^2 = x^2 + 16 - 8x$

⇒ $3x^2 + 8x + 4y^2 = 16$

⇒ $3\left(x + \frac{4}{3}\right)^2 + 4y^2 = \frac{64}{3}$

⇒ $\frac{(x + 4/3)^2}{64/9} + \frac{y^2}{16/3} = 1$

⇒ $a = 8/3$

21. If the straight lines $\frac{x-1}{k} = \frac{y-2}{2} = \frac{z-3}{3}$ and $\frac{x-2}{3} = \frac{y-3}{k} = \frac{z-1}{2}$ intersect at a point, then the integer k is equal to

- (1) 5 (2) 2 (3) -2 (4) -5

Sol: Ans [4]

$\frac{x-1}{k} = \frac{y-2}{2} = \frac{z-3}{3} = \lambda \dots(i)$

$$\frac{x-2}{3} = \frac{y-3}{k} = \frac{z-1}{2} = \mu \quad \dots(ii)$$

∴ Point on the line (i)

$$(\lambda k + 1, 2\lambda + 2, 3\lambda + 3)$$

Points will satisfy the equation (ii)

$$\Rightarrow \frac{\lambda k + 1 - 2}{3} = \frac{2\lambda + 2 - 3}{k} = \frac{2\lambda + 3 - 1}{2}$$

After solving

$$\Rightarrow k = -5$$

22. The line passing through the points $(5, 1, a)$ and $(3, b, 1)$ crosses the yz -plane at the point

$$\left(0, \frac{17}{2}, -\frac{13}{2}\right). \text{ Then}$$

- (1) $a = 4, b = 6$ (2) $a = 6, b = 4$ (3) $a = 8, b = 2$ (4) $a = 2, b = 8$

Sol: Ans [2]

Equation of line passing from two points $(5, 1, a)$ and $(3, b, 1)$

$$\Rightarrow \frac{x-5}{2} = \frac{y-1}{1-b} = \frac{z-a}{a-1} = k \text{ (at that point)}$$

$$\Rightarrow (x, y, z) = (2k + 5, (1-b)k + 1, (a-1)k + a)$$

$$\text{Crosses } yz \text{ plane at } \left(0, \frac{17}{2}, -\frac{13}{2}\right)$$

$$\Rightarrow 2k + 5 = 0 \quad \Rightarrow k = -\frac{5}{2}$$

$$\Rightarrow (1-b)k + 1 = \frac{17}{2} \quad \Rightarrow (1-b)\left(-\frac{5}{2}\right) = \frac{17}{2} - 1 = \frac{15}{2}$$

$$\Rightarrow 1-b = -3$$

$$\Rightarrow b = 4$$

$$\Rightarrow (a-1)k + a = -\frac{13}{2} \quad \Rightarrow (a-1)\left(-\frac{5}{2}\right) + a = -\frac{13}{2}$$

$$\Rightarrow -5(a-1) + 2a = -13$$

$$\Rightarrow -3a + 5 = -13$$

$$\Rightarrow -3a = -18$$

$$\Rightarrow a = 6$$

$$(a, b) = (6, 4)$$

23. The non-zero vectors \mathbf{a} , \mathbf{b} and \mathbf{c} are related by $\mathbf{a} = 8\mathbf{b}$ and $\mathbf{c} = -7\mathbf{b}$. Then the angle between \mathbf{a} and \mathbf{c} is

- (1) $\pi/4$ (2) $\pi/2$ (3) π (4) 0

Sol: Ans [3]

$$\mathbf{a} = 8\mathbf{b}, \mathbf{c} = -7\mathbf{b}$$

$$\therefore \mathbf{a} = -\frac{8}{7}\mathbf{c}$$

\Rightarrow \mathbf{a} and \mathbf{c} are collinear and opposite direction. So angle between them is π .

24. The vector $\mathbf{a} = \alpha\mathbf{i} + 2\mathbf{j} + \beta\mathbf{k}$ lies in the plane of the vectors $\mathbf{b} = \mathbf{i} + \mathbf{j}$ and $\mathbf{c} = \mathbf{j} + \mathbf{k}$ and bisects the angle between \mathbf{b} and \mathbf{c} . Then which one of the following gives possible values of α and β ?

- (1) $\alpha = 1, \beta = 2$ (2) $\alpha = 2, \beta = 1$ (3) $\alpha = 1, \beta = 1$ (4) $\alpha = 2, \beta = 2$

Sol: Ans [3]

$$\mathbf{a} = \alpha\mathbf{i} + 2\mathbf{j} + \beta\mathbf{k}; \mathbf{b} = \mathbf{i} + \mathbf{j}; \mathbf{c} = \mathbf{j} + \mathbf{k}$$

$$\text{as lie in a plane} \Rightarrow \begin{vmatrix} \alpha & 2 & \beta \\ 1 & 1 & 0 \\ 0 & 1 & 1 \end{vmatrix} = 0$$

$$\Rightarrow \alpha(1) - 2(1) + \beta(1) = 0$$

$$\Rightarrow \alpha + \beta = 2$$

\Rightarrow Only (3) is the possibility.

25. The mean of the numbers $a, b, 8, 5, 10$ is 6 and the variance is 6.80. Then which one of the following gives possible values of a and b ?

- (1) $a = 5, b = 2$ (2) $a = 1, b = 6$ (3) $a = 3, b = 4$ (4) $a = 0, b = 7$

Sol: Ans [3]

$$\text{Mean} = 6 = \frac{a + b + 8 + 5 + 10}{5}$$

$$a + b = 7 \quad \dots(i)$$

$$\text{Variance} = 6.8 = \frac{a^2 + b^2 + 189}{5} - 36$$

$$a^2 + b^2 = 25$$

$$\Rightarrow a = 3, b = 4$$

26. A die is thrown. Let A be the event that the number obtained is greater than 3. Let B be the event that the number obtained is less than 5. Then $P(A \cup B)$ is

- (1) 0 (2) 1 (3) $\frac{2}{5}$ (4) $\frac{3}{5}$

Sol: Ans [2]

$$A = \{x : x > 3\} = \{4, 5, 6\}$$

$$B = \{1, 2, 3, 4\}$$

$$P(A) = \frac{3}{6} = \frac{1}{2}; \quad P(B) = \frac{4}{6} = \frac{2}{3}$$

$$P(A \cap B) = \frac{1}{6}$$

$$\Rightarrow P(A \cup B) = \frac{1}{2} + \frac{2}{3} - \frac{1}{6} = \frac{3 + 4 - 1}{6} = 1$$

27. It is given that the events A and B are such that $P(A) = 1/4$, $P(A|B) = 1/2$ and $P(B|A) = 2/3$. Then $P(B)$ is
 (1) $1/3$ (2) $2/3$ (3) $1/2$ (4) $1/6$

Sol: Ans [1]

$$P(A) = 1/4, P(A|B) = 1/2 \text{ and } P(B|A) = 2/3$$

$$P(B) = ?$$

$$P\left(\frac{B}{A}\right) = \frac{P(B \cap A)}{P(A)}$$

$$\frac{2}{3} = \frac{P(B \cap A)}{1/4}$$

$$\Rightarrow P(B \cap A) = \frac{1}{6}$$

$$P\left(\frac{A}{B}\right) = \frac{P(A \cap B)}{P(B)} = \frac{1/6}{P(B)}$$

$$\frac{1}{2} = \frac{1}{6P(B)}$$

$$\Rightarrow P(B) = \frac{1}{3}$$

28. AB is a vertical pole with B at the ground level and A at the top. A man finds that the angel of elevation of the point A from a certain point C on the ground is 60° . He moves away from the pole along the line BC to a point D such that $CD = 7$ m. From D the angle of elevation of the point A is 45° . Then the height of the pole is

- (1) $\frac{7\sqrt{3}}{2}(\sqrt{3} + 1)$ m (2) $\frac{7\sqrt{3}}{2}(\sqrt{3} - 1)$ m (3) $\frac{7\sqrt{3}}{2} \frac{1}{\sqrt{3} + 1}$ m (4) $\frac{7\sqrt{3}}{2} \frac{1}{\sqrt{3} - 1}$ m

Sol: Ans [1]

$$\tan 60^\circ = \frac{h}{x} \Rightarrow x = \frac{h}{\sqrt{3}}$$


$$\tan 45^\circ = \frac{h}{x + 7} \Rightarrow h = x + 7$$

$$\Rightarrow h = \frac{h}{\sqrt{3}} + 7$$

$$\Rightarrow h \left(\frac{\sqrt{3} - 1}{\sqrt{3}} \right) = 7$$

$$\Rightarrow h = \frac{7\sqrt{3}}{\sqrt{3} - 1}$$

$$\Rightarrow h = \frac{7\sqrt{3}}{2}(\sqrt{3} + 1)$$



29. The value of $\cot\left(\operatorname{cosec}^{-1}\frac{5}{3} + \tan^{-1}\frac{2}{3}\right)$ is

- (1) $\frac{3}{17}$ (2) $\frac{4}{17}$ (3) $\frac{5}{17}$ (4) $\frac{6}{17}$

Sol: Ans [4]

$$\begin{aligned} \cot\left[\operatorname{cosec}^{-1}\frac{5}{3} + \cot^{-1}\frac{3}{2}\right] &= \cot\left[\cot^{-1}\frac{4}{3} + \cot^{-1}\frac{3}{2}\right] \\ &= \frac{2-1}{\frac{4}{3} + \frac{3}{2}} = \frac{1}{\frac{8+9}{6}} = \frac{6}{17} \end{aligned}$$

30. The statement $p \rightarrow (q \rightarrow p)$ is equivalent to

- (1) $p \rightarrow (p \vee q)$ (2) $p \rightarrow (p \wedge q)$ (3) $p \rightarrow (p \leftrightarrow q)$ (4) $p \rightarrow (p \rightarrow q)$

Sol: Ans [1]

p	q	$q \rightarrow p$	$p \rightarrow (q \rightarrow p)$	$p \vee q$	$p \rightarrow (p \vee q)$
0	0	1	1	0	1
0	1	0	1	1	1
1	0	1	1	1	1
1	1	1	1	1	1

Directions: Question number 31 to 35 are **Assertion-Reason** type questions. Each of these questions contains two statements: **Statement-1 (Assertion)** and **Statement-2 (Reason)**. Each of these questions also has four alternative choices, only one of which is the correct answer. You have to select the correct choice.

31. Let A be a 2×2 matrix with real entries. Let I be the 2×2 identity matrix. Denote by $\operatorname{tr}(A)$, the sum of diagonal entries of A . Assume that $A^2 = I$.

Statement-1: If $A \neq I$ and $A \neq -I$, then $\det A = -1$.

Statement-2: If $A \neq I$ and $A \neq -I$, then $\operatorname{tr}(A) \neq 0$.

- (1) Statement-1 is true, Statement-2 is true; Statement-2 is a correct explanation for Statement-1.
 (2) Statement-1 is true, Statement-2 is true; Statement-2 is not a correct explanation for Statement-1.
 (3) Statement-1 is true, Statement-2 is false.
 (4) Statement-1 is false, Statement-2 is true.

Sol: Ans [3]

Let $A = \begin{bmatrix} x_1 & x_2 \\ x_3 & x_4 \end{bmatrix}$ Again $A^2 = I$

$$\begin{aligned} \Rightarrow x_1^2 + x_2x_3 &= 1 & x_2(x_1 + x_4) &= 0 \\ x_4^2 + x_2x_3 &= 1 & x_3(x_1 + x_4) &= 0 \end{aligned}$$

Case-I: If $(x_1 + x_4) = 0 \Rightarrow \text{tr}(A) = 0$

Also it is not I and $-I$

Case-II: If $x_2 = 0 \Rightarrow x_3 = 0 \Rightarrow x_1 = \pm 1$ and $x_4 = \pm 1$

There are two cases $A = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$ & $\begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix}$

For which $\det A = -1$

Both cases describe answer is (3)

32. **Statement-1:** For every natural number $n \geq 2$, $\frac{1}{\sqrt{1}} + \frac{1}{\sqrt{2}} + \dots + \frac{1}{\sqrt{n}} > \sqrt{n}$.

Statement-2: For every natural number $n \geq 2$, $\sqrt{n(n+1)} < n+1$.

(1) Statement-1 is true, Statement-2 is true; Statement-2 is a correct explanation for Statement-1.

(2) Statement-1 is true, Statement-2 is true; Statement-2 is not a correct explanation for Statement-1.

(3) Statement-1 is true, Statement-2 is false.

(4) Statement-1 is false, Statement-2 is true.

Sol: Ans [2]

$$\begin{aligned} \frac{1}{\sqrt{1}} + \frac{1}{\sqrt{2}} + \frac{1}{\sqrt{3}} + \dots + \frac{1}{\sqrt{n}} + \frac{1}{\sqrt{n+1}} &> \sqrt{n} + \frac{1}{\sqrt{n+1}} \\ &> \frac{\sqrt{n(n+1)} + 1}{\sqrt{n+1}} > \sqrt{n+1} \end{aligned}$$

Since $\frac{\sqrt{n(n+1)} + 1}{\sqrt{n+1}} - \sqrt{n+1} > 0 \Rightarrow$ Statement-1 is true.

Again $\sqrt{n(n+1)} < (n+1)$

$\Rightarrow \sqrt{n} < \sqrt{n+1}$, which is true \Rightarrow (2) is answer

33. **Statement-1:** $\sum_{r=0}^n (r+1) {}^n C_r = (n+2)2^{n-1}$.

Statement-2: $\sum_{r=0}^n (r+1) {}^n C_r x^r = (1+x)^n + nx(1+x)^{n-1}$.

(1) Statement-1 is true, Statement-2 is true; Statement-2 is a correct explanation for Statement-1.

(2) Statement-1 is true, Statement-2 is true; Statement-2 is not a correct explanation for Statement-1.

(3) Statement-1 is true, Statement-2 is false.

(4) Statement-1 is false, Statement-2 is true.

Sol: Ans [1]

Since $\sum_{r=0}^n (r+1) {}^n C_r x^r = (1+x)^n + nx(1+x)^{n-1}$

Putting $x = 1$, we get $\sum_{r=0}^n (r+1) {}^n C_r = (n+2)2^{n-1}$

\Rightarrow (1) is answer

34. In a shop there are five types of ice-creams available. A child buys six ice-creams.

Statement-1: The number of different ways the child can buy the six ice-creams is ${}^{10}C_5$.

Statement-2: The number of different ways the child can buy the six ice-creams is equal to the number of different ways of arranging 6 A's and 4 B's in a row.

(1) Statement-1 is true, Statement-2 is true; Statement-2 is a correct explanation for Statement-1.

(2) Statement-1 is true, Statement-2 is true; Statement-2 is not a correct explanation for Statement-1.

(3) Statement-1 is true, Statement-2 is false.

(4) Statement-1 is false, Statement-2 is true.

Sol: Ans [4]

The no. of ways = ${}^{6+5-1}C_6 = {}^{10}C_6$

No. of ways of arranging 6 A's and 4 B's in a row = $\frac{10!}{6!4!} = {}^{10}C_6$

\Rightarrow Statement-1 is false and statement-2 is true.

35. Let p be the statement "x is an irrational number", q be the statement "y is a transcendental number" and r be the statement "x is a rational number iff y is a transcendental number".

Statement-1: r is equivalent to either q or p .

Statement-2: r is equivalent to $\sim(p \leftrightarrow \sim q)$.

(1) Statement-1 is true, Statement-2 is true; Statement-2 is a correct explanation for Statement-1.

(2) Statement-1 is true, Statement-2 is true; Statement-2 is not a correct explanation for Statement-1.

(3) Statement-1 is true, Statement-2 is false.

(4) Statement-1 is false, Statement-2 is true.

Sol: Ans []

Clearly r is equivalent to $\sim p \leftrightarrow q$

p	q	$\sim p$	r	q or p	$\sim q$	$p \leftrightarrow \sim q$	$\sim(p \leftrightarrow \sim q)$
0	0	1	0	0	1	0	1
0	1	1	1	1	0	1	0
1	0	0	1	1	1	1	0
1	1	0	0	1	0	0	1

Clearly r is neither equivalent to q or p nor to $\sim(p \leftrightarrow \sim q)$

So statement-1 and 2 both are false.

PART-B: CHEMISTRY

36. The ionization enthalpy of hydrogen atom is $1.312 \times 10^6 \text{ J mol}^{-1}$. The energy required to excite the electron in the atom from $n = 1$ to $n = 2$ is

- (1) $6.56 \times 10^5 \text{ J mol}^{-1}$ (2) $7.56 \times 10^5 \text{ J mol}^{-1}$ (3) $9.84 \times 10^5 \text{ J mol}^{-1}$ (4) $8.51 \times 10^5 \text{ J mol}^{-1}$

Sol: Ans [3] $E_1 = -1.312 \times 10^6 \text{ J/mol}$

$$\Delta E = E_2 - E_1$$

37. Which one of the following pairs of species have the same bond order?

- (1) CN^- and CN^+ (2) O_2^- and CN^- (3) NO^+ and CN^+ (4) CN^- and NO^+

Sol: Ans [4] Both CN^- and NO^+ have bond order = 3.

38. Which of the following constitutes a group of the isoelectronic species?

- (1) NO^+ , C_2^{2-} , CN^- , N_2 (2) CN^- , N_2 , O_2^{2-} , C_2^{2-} (3) N_2 , O_2^- , NO^+ , CO (4) C_2^{2-} , O_2^- , CO , NO

Sol: Ans [1]

39. Four species are listed below:

- (i) HCO_3^- (ii) H_3O^+ (iii) HSO_4^- (iv) HSO_3F

Which one of the following is the correct sequence of the acid strength?

- (1) $\text{ii} < \text{iii} < \text{i} < \text{iv}$ (2) $\text{i} < \text{iii} < \text{ii} < \text{iv}$ (3) $\text{iii} < \text{i} < \text{iv} < \text{ii}$ (4) $\text{iv} < \text{ii} < \text{iii} < \text{i}$

Sol: Ans [2]

40. The pK_a of a weak acid, HA, is 4.80. The pK_b of a weak base, BOH, is 4.78. The pH of an aqueous solution of the corresponding salt, BA, will be

- (1) 4.79 (2) 7.01 (3) 9.22 (4) 9.58

Sol: Ans [2] $\text{pH} = \frac{1}{2} (\text{pK}_w + \text{pK}_a - \text{pK}_b)$

41. The correct decreasing order of priority for the functional groups of organic compounds in the IUPAC system of nomenclature is

- (1) $-\text{SO}_3\text{H}$, $-\text{COOH}$, $-\text{CONH}_2$, $-\text{CHO}$ (2) $-\text{CHO}$, $-\text{COOH}$, $-\text{SO}_3\text{H}$, $-\text{CONH}_2$
 (3) $-\text{CONH}_2$, $-\text{CHO}$, $-\text{SO}_3\text{H}$, $-\text{COOH}$ (4) $-\text{COOH}$, $-\text{SO}_3\text{H}$, $-\text{CONH}_2$, $-\text{CHO}$

Sol: Ans [1]

42. The treatment of CH_3MgX with $\text{CH}_3\text{C} \equiv \text{C} - \text{H}$ produces

- (1) $\text{CH}_3\text{C} \equiv \text{C} - \text{CH}_3$ (2) $\text{CH}_3 - \overset{\text{H}}{\underset{|}{\text{C}}} = \overset{\text{H}}{\underset{|}{\text{C}}} - \text{CH}_3$ (3) CH_4 (4) $\text{CH}_3 - \text{CH} = \text{CH}_2$

Sol: Ans [3] $\text{R-MgX} + \text{compound having acidic hydrogen} \rightarrow \text{RH}$

43. The hydrocarbon which can react with sodium in liquid ammonia is

- (1) $\text{CH}_3\text{CH}_2\text{C} \equiv \text{CH}$
- (2) $\text{CH}_3\text{CH} = \text{CHCH}_3$
- (3) $\text{CH}_3\text{CH}_2\text{C} \equiv \text{CCH}_2\text{CH}_3$
- (4) $\text{CH}_3\text{CH}_2\text{CH}_2\text{C} \equiv \text{CCH}_2\text{CH}_2\text{CH}_3$

Sol: Ans [1]

44. The vapour pressure of water at 20°C is 17.5 mm Hg. If 18 g of glucose ($\text{C}_6\text{H}_{12}\text{O}_6$) is added to 178.2 g of water at 20°C , the vapour pressure of the resulting solution will be

- (1) 15.750 mm Hg
- (2) 16.500 mm Hg
- (3) 17.325 mm Hg
- (4) 17.675 mm Hg.

Sol: Ans [3] $\frac{P_A^0 - P_A}{P_A^0} = x_B$

45. Gold numbers of protective colloids A, B, C and D are 0.50, 0.01, 0.10 and 0.005, respectively. The correct order of their protective powers is

- (1) $\text{C} < \text{B} < \text{D} < \text{A}$
- (2) $\text{A} < \text{C} < \text{B} < \text{D}$
- (3) $\text{B} < \text{D} < \text{A} < \text{C}$
- (4) $\text{D} < \text{A} < \text{C} < \text{B}$

Sol: Ans [2] More the gold number lesser is the protecting power.

46. In a compound, atoms of element Y form ccp lattice and those of element X occupy $2/3^{\text{rd}}$ of tetrahedral voids. The formula of the compound will be

- (1) X_2Y_3
- (2) X_2Y
- (3) X_3Y_4
- (4) X_4Y_3

Sol: Ans [4] atoms Y = 4, $\text{X} = 8 \times \frac{2}{3} = \frac{16}{3}$

$$\text{hence } \text{X}_{16/3} \text{Y}_4 = \text{X}_4\text{Y}_3$$

47. In context with the industrial preparation of hydrogen from water gas ($\text{CO} + \text{H}_2$), which of the following is the correct statement?

- (1) CO is removed by absorption in aqueous Cu_2Cl_2 solution
- (2) H_2 is removed through occlusion with Pd
- (3) CO is oxidised to CO_2 with steam in the presence of a catalyst followed by absorption of CO_2 in alkali
- (4) CO and H_2 are fractionally separated using differences in their densities

Sol: Ans [3]

48. Among the following substituted silanes the one which will give rise to cross linked silicone polymer on hydrolysis is

- (1) RSiCl_3
- (2) R_2SiCl_2
- (3) R_3SiCl
- (4) R_4Si

Sol: Ans [1] RSiCl_3 on hydrolysis give RSi(OH)_3 which generate cross links, by polymerisation.

49. Amount of oxalic acid present in a solution can be determined by its titration with KMnO_4 solution in the presence of H_2SO_4 . The titration gives unsatisfactory result when carried out in the presence of HCl because HCl

- (1) furnishes H^+ ions in addition to those from oxalic acid
- (2) reduces permanganate to Mn^{2+}
- (3) oxidises oxalic acid to carbon dioxide and water
- (4) gets oxidised by oxalic acid to chlorine

Sol: Ans [2] HCl reacts with KMnO_4 to form MnCl_2 and Cl_2

50. Given $E_{\text{Cr}^{3+}/\text{Cr}}^\circ = -0.72 \text{ V}$, $E_{\text{Fe}^{2+}/\text{Fe}}^\circ = -0.42 \text{ V}$. The potential for the cell

$\text{Cr}|\text{Cr}^{3+} (0.1 \text{ M})||\text{Fe}^{2+} (0.01 \text{ M})|\text{Fe}$ is

- (1) 0.339 V
- (2) -0.339 V
- (3) -0.26 V
- (4) 0.26 V

Sol: Ans [4] $E_{\text{cell}} = E_{\text{cell}}^\circ - \frac{0.0591}{6} \log \frac{[\text{Cr}^{3+}]^2}{[\text{Fe}^{2+}]^3}$.

51. Which one of the following is the correct statement?

- (1) Beryllium exhibits coordination number of six
- (2) Chlorides of both beryllium and aluminium have bridged chloride structures in solid phase
- (3) $\text{B}_2\text{H}_6 \cdot 2\text{NH}_3$ is known as 'inorganic benzene'
- (4) Boric acid is a protonic acid

Sol: Ans [2]

52. Identify the **wrong** statement in the following:

- (1) Greenhouse effect is responsible for global warming
- (2) Ozone layer does not permit infrared radiation from the sun to reach the earth
- (3) Acid rain is mostly because of oxides of nitrogen and sulphur
- (4) Chlorofluorocarbons are responsible for ozone layer depletion

Sol: Ans [2]

53. The coordination number and the oxidation state of the element 'E' in the complex $[\text{E}(\text{en})_2(\text{C}_2\text{O}_4)]\text{NO}_2$ (when (en) is ethylene diamine) are, respectively

- (1) 4 and 2
- (2) 4 and 3
- (3) 6 and 3
- (4) 6 and 2

Sol: Ans [3] en and $\text{C}_2\text{O}_4^{2-}$ are both bidentate ligands.

54. In which of the following octahedral complexes of Co (at. no. 27), will the magnitude of Δ_0 be the highest?

- (1) $[\text{Co}(\text{C}_2\text{O}_4)_3]^{3-}$
- (2) $[\text{Co}(\text{H}_2\text{O})_6]^{3+}$
- (3) $[\text{Co}(\text{NH}_3)_6]^{3+}$
- (4) $[\text{Co}(\text{CN})_6]^{3-}$

Sol: Ans [4]

55. Larger number of oxidation states are exhibited by the actinoids than those by the lanthanoids, the main reason being

- (1) lesser energy difference between 5f and 6d than between 4f and 5d orbitals
- (2) more energy difference between 5f and 6d than between 4f and 5d orbitals
- (3) more reactive nature of the actinoids than the lanthanoids
- (4) 4f orbitals are more diffused than the 5f orbitals


Sol: Ans [1]

56. Which of the following factors is of *no significance* for roasting sulphide ores to the oxides and not subjecting the sulphide ores to carbon reduction directly?


- (1) CO_2 is thermodynamically more stable than CS_2
- (2) Metal sulphides are less stable than the corresponding oxides
- (3) CO_2 is more volatile than CS_2
- (4) Metal sulphides are thermodynamically more stable than CS_2

Sol: Ans [4]

57. Oxidising power of chlorine in aqueous solution can be determined by the parameters indicated below



The energy involved in the conversion of




(using the data $\Delta_{\text{diss}} \text{H}_{\text{Cl}_2}^\ominus = 240 \text{ kJ mol}^{-1}$, $\Delta_{\text{eg}} \text{H}_{\text{Cl}}^\ominus = -349 \text{ kJ mol}^{-1}$, $\Delta_{\text{hyd}} \text{H}_{\text{Cl}^\ominus}^\ominus = -381 \text{ kJ mol}^{-1}$) will be

- (1) -610 kJ mol^{-1}
- (2) -850 kJ mol^{-1}
- (3) $+120 \text{ kJ mol}^{-1}$
- (4) $+152 \text{ kJ mol}^{-1}$

Sol: Ans [1] $\Delta H = \Delta H_1 + \Delta H_2 + \Delta H_3 = \frac{1}{2} \times 240 - 349 - 381 = -610 \text{ kJ/mol}$

58. In the following sequence of reactions, the alkene affords the compound 'B'



- (1) CH_3COCH_3
- (2) $\text{CH}_3\text{CH}_2\text{COCH}_3$
- (3) CH_3CHO
- (4) $\text{CH}_3\text{CH}_2\text{CHO}$

Sol: Ans [3]

59. Phenol, when it first reacts with concentrated sulphuric acid and then with concentrated nitric acid, gives

- (1) *o*-nitrophenol
- (2) *p*-nitrophenol
- (3) nitrobenzene
- (4) 2,4,6-trinitrobenzene

Sol: Ans [1]

60. Toluene is nitrated and the resulting product is reduced with tin and hydrochloric acid. The product so obtained is diazotised and then heated with cuprous bromide. The reaction mixture so formed contains


- (1) mixture of *o*- and *p*-dibromobenzenes (2) mixture of *o*- and *p*-bromoanilines
 (3) mixture of *o*- and *m*-bromotoluenes (4) mixture of *o*- and *p*-bromotoluenes

Sol: Ans [4] $\text{Ph-CH}_3 \xrightarrow[\text{H}_2\text{SO}_4]{\text{HNO}_3} \text{o, p-nitrotoluene} \xrightarrow{\text{Sn/HCl}} \text{o, p-methyl aniline}$
 $\xrightarrow[0^\circ\text{C}]{\text{NaNO}_2/\text{HCl}} \text{o, p-methyl benzene diazonium chloride} \xrightarrow{\text{CuBr}} \text{o, p-bromotoluene.}$

61. The organic chloro compound, which shows complete stereochemical inversion during a $\text{S}_{\text{N}}2$ reaction, is

- (1) $(\text{CH}_3)_3\text{CCl}$ (2) $(\text{CH}_3)_2\text{CHCl}$ (3) CH_3Cl (4) $(\text{C}_2\text{H}_5)_2\text{CHCl}$

Sol: Ans [3] Lesser the steric hindrance, more is the possibility of $\text{S}_{\text{N}}2$

62. The absolute configuration of  is

- (1) R, R (2) R, S (3) S, R (4) S, S


Sol: Ans [1]

63. α -D-(+)-glucose and β -D-(+)-glucose are

- (1) epimers (2) anomers (3) enantiomers (4) conformers


Sol: Ans [2]

64. The electrophile, E^\oplus attacks the benzene ring to generate the intermediate σ -complex. Of the following, which σ -complex is of lowest energy?



Sol: Ans [1]




65. Standard entropy of X_2 , Y_2 and XY_3 are 60, 40 and 50 $\text{JK}^{-1} \text{mol}^{-1}$, respectively. For the reaction,



- (1) 500 K (2) 750 K (3) 1000 K (4) 1250 K

Sol: Ans [2] $\Delta S = \frac{?H}{T}$ or $T = \frac{?H}{?S} = \frac{-30 \times 1000}{50 - \frac{1}{2} \times 60 - \frac{3}{2} \times 40} = 750 \text{ K}$

66. For the following three reactions a, b and c equilibrium constants are given:



Which of the following relation is correct?

(1) $K_2K_3 = K_1$ (2) $K_3 = K_1K_2$ (3) $K_3 \cdot K_2^3 = K_1^2$ (4) $K_1\sqrt{K_2} = K_3$

Sol: Ans [2]

67. Bakelite is obtained from phenol by reacting with



Sol: Ans [3]

68. The equilibrium constants K_{P_1} and K_{P_2} for the reactions $\text{X} \rightleftharpoons 2\text{Y}$ and $\text{Z} \rightleftharpoons \text{P} + \text{Q}$, respectively are in the ratio of 1 : 9. If the degree of dissociation of X and Z be equal then the ratio of total pressures at these equilibria is

(1) 1 : 1 (2) 1 : 3 (3) 1 : 9 (4) 1 : 36

Sol: Ans [4]

69. For a reaction $\frac{1}{2}\text{A} \rightarrow 2\text{B}$, rate of disappearance of 'A' is related to the rate of appearance of 'B' by the expression

(1) $-\frac{d[\text{A}]}{dt} = \frac{1}{4} \frac{d[\text{B}]}{dt}$ (2) $-\frac{d[\text{A}]}{dt} = \frac{d[\text{B}]}{dt}$

(3) $-\frac{d[\text{A}]}{dt} = 4 \frac{d[\text{B}]}{dt}$ (4) $-\frac{d[\text{A}]}{dt} = \frac{1}{2} \frac{d[\text{B}]}{dt}$

Sol: Ans [1] Rate of reaction = $\frac{-d[\text{A}]}{\frac{1}{2}dt} = \frac{dB}{2dt}$ or $\frac{-d[\text{A}]}{dt} = \frac{1}{4} \frac{dB}{dt}$

70. At 80°C, the vapour pressure of pure liquid 'A' is 520 mm Hg and that of pure liquid 'B' is 1000 mm Hg. If a mixture solution of 'A' and 'B' boils at 80°C and 1 atm pressure, the amount of 'A' in the mixture is (1 atm = 760 mm Hg)

(1) 34 mol percent (2) 48 mol percent (3) 50 mol percent (4) 52 mol percent

Sol: Ans [3] $P = P_A^\circ x_A + P_B^\circ x_B = P_A^\circ x_A + P_B^\circ (1 - x_A)$

$x_A = 0.5.$

Percentage = $0.5 \times 100 = 50\%$

PART-C: PHYSICS

71. A body of mass $m = 3.513$ kg is moving along the x -axis with a speed of 5.00 ms^{-1} . The magnitude of its momentum is recorded as

- (1) 17.565 kg ms^{-1} (2) 17.55 kg ms^{-1} (3) 17.57 kg ms^{-1} (4) 17.6 kg ms^{-1}

Sol: Ans [4] $p = mv = 3.513 \times 5.00 = 17.565 = 17.6$

72. Consider a uniform square plate of side a and mass m . The moment of inertia of this plate about an axis perpendicular to its plane and passing through one of its corners is

- (1) $\frac{1}{12}ma^2$ (2) $\frac{7}{12}ma^2$ (3) $\frac{2}{3}ma^2$ (4) $\frac{5}{6}ma^2$

Sol: Ans [3] $I_z = I_x + I_y = \frac{ma^2}{3} + \frac{ma^2}{3} = \frac{2}{3}ma^2$

73. The speed of sound in oxygen (O_2) at a certain temperature is 460 ms^{-1} . The speed of sound in helium (He) at the same temperature will be (assume both gases to be ideal)


- (1) 500 ms^{-1} (2) 650 ms^{-1} (3) 330 ms^{-1} (4) 460 ms^{-1}

Sol: Ans [not matching] $460 = \sqrt{\frac{\frac{7}{5}RT}{32}}$

$$v = \sqrt{\frac{\frac{5}{3}RT}{4}}$$

$$\therefore v = 460 \sqrt{\frac{32}{4} \times \frac{5}{3} \times \frac{5}{7}} \approx 1462 \text{ m/s}$$

74. A thin spherical shell of radius R has charge Q spread uniformly over its surface. Which of the following graphs most closely represents the electric field $E(r)$ produced by the shell in the range $0 \leq r < \infty$, where r is the distance from the centre of the shell ?



Sol: Ans [4] $E_{\text{in}} = 0$

$$E_{\text{out}} = K \frac{Q}{r^2}$$

75. Relative permittivity and permeability of a material are ϵ_r and μ_r , respectively. Which of the following values of these quantities are allowed for a diamagnetic material ?

- (1) $\epsilon_r = 1.5, \mu_r = 0.5$ (2) $\epsilon_r = 0.5, \mu_r = 0.5$ (3) $\epsilon_r = 1.5, \mu_r = 1.5$ (4) $\epsilon_r = 0.5, \mu_r = 1.5$

Sol: Ans [1] For diamagnetic material $\mu_r < 1$ and $\epsilon_r > 1$

76. Suppose an electron is attracted towards the origin by a force $\frac{k}{r}$ where k is a constant and r is the distance of the electron from the origin. By applying Bohr model to this system, the radius of the n^{th} orbital of the electron is found to be r_n and the kinetic energy of the electron to be T_n . Then which of the following is true ?

- (1) T_n independent of n , $r_n \propto n$ (2) $T_n \propto \frac{1}{n}$, $r_n \propto n$
 (3) $T_n \propto \frac{1}{n}$, $r_n \propto n^2$ (4) $T_n \propto \frac{1}{n^2}$, $r_n \propto n^2$

Sol: Ans [1] $m \frac{v^2}{r} = \frac{k}{r}$

$\Rightarrow mv^2 = k$

$\therefore T_n = \frac{1}{2}mv^2 = \frac{k}{2}$

$\Rightarrow T_n$ independent of n .

77. A block of mass 0.50 kg is moving with a speed of 2.00 ms⁻¹ on a smooth surface. It strikes another mass of 1.00 kg and then they move together as a single body. The energy loss during the collision is

- (1) 1.00 J (2) 0.67 J (3) 0.34 J (4) 0.16 J

Sol: Ans [2] Loss = $\frac{1}{2} \times \frac{0.5 \times 1}{0.5+1} 2^2 (1-0^2) = 2 \times \frac{0.5}{1.5} = \frac{2}{3} J$

78. A wave travelling along the x -axis is described by the equation $y(x, t) = 0.005 \cos (\alpha x - \beta t)$. If the wavelength and the time period of the wave are 0.08 m and 2.0 s, respectively, then α and β in appropriate units are

- (1) $\alpha = \frac{0.08}{\pi}, \beta = \frac{2.0}{\pi}$ (2) $\alpha = \frac{0.04}{\pi}, \beta = \frac{1.0}{\pi}$ (3) $\alpha = 12.50\pi, \beta = \frac{\pi}{2.0}$ (4) $\alpha = 25.00 \pi, \beta = \pi$

Sol: Ans [4] $\alpha = \frac{2\pi}{0.08} = 25\pi$


$\beta = \frac{2\pi}{2} = \pi$

79. A working transistor with its three legs marked P , Q and R is tested using a multimeter. No conduction is found between P and Q . By connecting the common (negative) terminal of the multimeter to R the other (positive) terminal to P or Q , some resistance is seen on the multimeter. Which of the following is true for the transistor ?

- (1) It is a pnp transistor with R as collector (2) It is a pnp transistor with R as emitter
 (3) It is an nnp transistor with R as collector (4) It is an nnp transistor with R as base

Sol: Ans [4] P and Q are same material because there is no conduction current even if polarity of multimeter is changed. So R should be base.

80. A jar is filled with two non-mixing liquids 1 and 2 having densities ρ_1 and ρ_2 , respectively. A solid ball, made up of a material of density ρ_3 , is dropped in the jar. It comes to equilibrium in the position shown in the figure



Which of the following is true for ρ_1 , ρ_2 and ρ_3 ?

- (1) $\rho_1 > \rho_3 > \rho_2$ (2) $\rho_1 < \rho_2 < \rho_3$
 (3) $\rho_1 < \rho_3 < \rho_2$ (4) $\rho_3 < \rho_1 < \rho_2$

Sol: Ans [3] Since ball dipped in liquid-1 $\Rightarrow \rho_3 > \rho_1$
 Ball floats in liquid-2 $\Rightarrow \rho_3 < \rho_2$

81. An athlete in the Olympic games covers a distance of 100 m in 10 s. His kinetic energy can be estimated to be in the range
- (1) $2 \times 10^5 \text{ J} - 3 \times 10^5 \text{ J}$ (2) 20,000 J – 50,000 J
 (3) 2,000 J – 5,000 J (4) 200 J – 500 J

Sol: Ans [3] $k_e = \frac{1}{2}mv^2$, take mass $\approx 60 \text{ kg} = \frac{1}{2} \times 60 \times (10)^2 = 3000$

82. A parallel plate capacitor with air between the plates has a capacitance of 9 pF. The separation between its plates is d . The space between the plates is now filled with two dielectrics. One of the dielectrics has dielectric constant $k_1 = 3$ and thickness $\frac{d}{3}$ while the other one has dielectric constant

$k_2 = 6$ and thickness $\frac{2d}{3}$. Capacitance of the capacitor is now

- (1) 45 pF (2) 40.5 pF (3) 20.25 pF (4) 1.8 pF

Sol: Ans [2] $\frac{\epsilon_0 A}{d} = 9 \text{ pF}$


$$C = \frac{\epsilon_0 A}{\frac{d}{3k_1} + \frac{2d}{3k_2}} = \frac{\epsilon_0 A}{d \left(\frac{1}{9} + \frac{2}{18} \right)} = \frac{9}{\left(\frac{1}{9} + \frac{2}{18} \right)} = 40.5 \text{ pF}$$

83. The dimension of magnetic field in M, L, T and C (Coulomb) is given as

- (1) $M T^2 C^{-2}$ (2) $M T^{-1} C^{-1}$ (3) $M T^{-2} C^{-1}$ (4) $M L T^{-1} C^{-1}$

Sol: Ans [2] $F = qvB \Rightarrow B = \frac{F}{qv} = \frac{MLT^{-2}}{CLT^{-1}} = MT^{-1}C^{-1}$

84. A body is at rest at $x = 0$. At $t = 0$, it starts moving in the positive x -direction with a constant acceleration. At the same instant another body passes through $x = 0$ moving in the positive x -direction with a constant speed. The position of the first body is given by $x_1(t)$ after time t and that of the second body by $x_2(t)$ after the same time interval. Which of the following graphs correctly describes $(x_1 - x_2)$ as a function of time t ?



Sol: Ans [1] $x_1(t) = \frac{1}{2}at^2$

$$x_2(t) = vt$$

$$x_1 - x_2 = \frac{1}{2}at^2 - vt = t\left(\frac{a}{2}t - v\right)$$


at $t = 0$, $x_1 - x_2 = 0$

at $t = \frac{2v}{a}$, $x_1 - x_2 = 0$

after $t = \frac{2v}{a}$, $x_1 - x_2$ increases

at $0 < t < \frac{2v}{a}$, $x_1 - x_2$ is negative.

85. In the circuit below, A and B represent two inputs and C represents the product



The circuit represents

- (1) AND gate (2) NAND gate (3) OR gate (4) NOR gate

Sol: Ans [3] Its truth table is

A	B	C
0	1	1
1	0	1
1	1	1
0	0	0

⇒ It is OR gate

86. While measuring the speed of sound by performing a resonance column experiment, a student gets the first resonance condition at a column length of 18 cm during winter. Repeating the same experiment during summer, she measures the column length to be x cm for the second resonance. Then

- (1) $x > 54$ (2) $54 > x > 36$ (3) $36 > x > 18$ (4) $18 > x$

Sol: Ans [1] $\frac{\lambda_w}{4} = 18 \text{ cm}$

$\lambda_w = 72 \text{ cm}$


For second resonance $\frac{3\lambda_s}{4} = x$

as velocity increased in summer $\Rightarrow \lambda_s > \lambda_w$

$\Rightarrow x > \frac{3 \times 72}{4}$

$x > 54 \text{ cm.}$

87. Shown in the figure below is a meter-bridge set up with null deflection in the galvanometer.



The value of the unknown resistor R is

- (1) 220 Ω (2) 110 Ω (3) 55 Ω (4) 13.75 Ω

Sol: Ans [1]

$\frac{55}{20} = \frac{R}{100 - 20} \Rightarrow R = 55 \times 4 = 220 \Omega$

88. A spherical solid ball of volume V is made of a material of density ρ_1 . It is falling through a liquid of density ρ_2 ($\rho_2 < \rho_1$). Assume that the liquid applies a viscous force on the ball that is proportional to the square of its speed v , i.e., $F_{\text{viscous}} = -kv^2$ ($k > 0$). The terminal speed of the ball is

- (1) $\frac{Vg\rho_1}{k}$ (2) $\sqrt{\frac{Vg\rho_1}{k}}$
 (3) $\frac{Vg(\rho_1 - \rho_2)}{k}$ (4) $\sqrt{\frac{Vg(\rho_1 - \rho_2)}{k}}$

Sol: Ans [4] At terminal speed, $kv_T^2 + \rho_2 Vg = \rho_1 Vg \Rightarrow v_T = \sqrt{\frac{(\rho_1 - \rho_2)Vg}{k}}$

Sol: Ans [4]

$$\begin{aligned}
 n_1 C_v T_1 + n_2 C_v T_2 &= (n_1 + n_2) C_v T \\
 \Rightarrow n_1 T_1 + n_2 T_2 &= (n_1 + n_2) T \\
 \frac{P_1 V_1}{RT_1} T_1 + \frac{P_2 V_2}{RT_2} T_2 &= \left(\frac{P_1 V_1}{RT_1} + \frac{P_2 V_2}{RT_2} \right) T \\
 P_1 V_1 + P_2 V_2 &= \left(\frac{P_1 V_1}{T_1} + \frac{P_2 V_2}{T_2} \right) T \\
 \therefore T &= \frac{(P_1 V_1 + P_2 V_2)(T_1 T_2)}{P_1 V_1 T_2 + P_2 V_2 T_1}
 \end{aligned}$$

- 92.** Two full turns of a circular scale of a screw gauge cover a distance of 1 mm on its main scale. The total number of divisions on the circular scale is 50. Further, it is found that the screw gauge has a zero error of -0.03 mm. While measuring the diameter of a thin wire, a student notes the main scale reading of 3 mm and the number of circular scale divisions in line with the main scale as 35. The diameter of the wire is
- (1) 3.73 mm (2) 3.67 mm (3) 3.38 mm (4) 3.32 mm

Sol: Ans [3] Least count = $\frac{(1/2)}{50} = \frac{1}{100}$ mm

Noted reading = $3 + 35 \times \frac{1}{100} = 3.35$ mm

\therefore Diameter = Reading + zero error = $3.35 + 0.03 = 3.38$ mm

- 93.** A horizontal overhead powerline is at a height of 4 m from the ground and carries a current of 100 A from east to west. The magnetic field directly below it on the ground is ($\mu_0 = 4\pi \times 10^{-7}$ T mA⁻¹)
- (1) 5×10^{-6} T northward (2) 5×10^{-6} T southward
 (3) 2.5×10^{-7} T northward (4) 2.5×10^{-7} southward


Sol: Ans [2] $B = \frac{\mu_0 I}{2\pi(4)} = \frac{\mu_0}{4\pi} \times \left(\frac{2I}{4} \right) = 10^{-7} \times \frac{2 \times 100}{4} = 50 \times 10^{-7}$

$\Rightarrow 5 \times 10^{-6}$ south

- 94.** An experiment is performed to find the refractive index of glass using a travelling microscope. In this experiment distance are measured by
- (1) a standard laboratory scale
 (2) a meter scale provided on the microscope
 (3) a screw gauge provided on the microscope
 (4) a vernier scale provided on the microscope

Sol: Ans [4] We need liner scale with high precision.

95. A 5 V battery with internal resistance 2Ω and 2 V battery with internal resistance 1Ω and is connected to a 10Ω resistor as shown in the figure




- (1) 0.03 A P_1 to P_2 (2) 0.03 A P_2 to P_1 (3) 0.27 A P_1 to P_2 (4) 0.27 A P_2 to P_1

Sol: Ans [2] Equivalent emf = $\frac{\frac{5}{2} - \frac{2}{1}}{\frac{1}{2} + \frac{1}{1}} = \frac{1}{3} V$

$$I = \frac{1/3}{10 + 2/3} = 0.03 A$$

96. A capillary tube (A) is dipped in water. Another identical tube (B) is dipped in a soap-water solution. Which of the following shows the relative nature of the liquid columns in the two tubes ?




Sol: Ans [2] Because surface tension reduces.

97. Two coaxial solenoids are made by winding thin insulated wire over a pipe of cross-sectional area $A = 10 \text{ cm}^2$ and length = 20 cm. If one of the solenoids has 300 turns and the other 400 turns, their mutual inductance is ($\mu_0 = 4\pi \times 10^{-7} \text{ T m A}^{-1}$)

- (1) $4.8\pi \times 10^{-4} \text{ H}$ (2) $4.8\pi \times 10^{-5} \text{ H}$ (3) $2.4\pi \times 10^{-4} \text{ H}$ (4) $2.4\pi \times 10^{-5} \text{ H}$

Sol: Ans [3] $M = \frac{\mu_0 N_1 N_2 A}{l} = \frac{4\pi \times 10^{-7} \times 300 \times 400 \times 10 \times 10^{-4}}{0.2} = 2.4\pi \times 10^{-4} \text{ H}$

98. A student measures the focal length of a convex lens by putting an object pin at a distance u from the lens and measuring the distance v of the image pin. The graph between u and v plotted by the student should look like



Sol: Ans [2] For real image u and v are of different sign and as $|u|$ increases $|v|$ decreases.

99. This question contains Statement-1 and Statement-2. Of the four choices given after the statements, choose the one that best describes the two statements.

Statement-1: For a mass M kept at the centre of a cube of side a , the flux of gravitational field passing through its sides is $4 \pi G M$

and

Statement-2: If the direction of a field due to a point source is radial and its dependence on the distance r from the source is given as $\frac{1}{r^2}$, its flux through a closed surface depends only on the strength of the source enclosed by the surface and not on the size or shape of the surface.

- (1) Statement-1 is true, Statement-2 is true; Statement-2 is a correct explanation for Statement-1
 (2) Statement-1 is true, Statement-2 is true; Statement-2 is not a correct explanation for Statement-1
 (3) Statement-1 is true, Statement-2 is false
 (4) Statement-1 is false, Statement-2 is true

Sol: Ans [1] Statement-1 is true based upon Gauss's theorem. Gauss's theorem is also based upon inverse square law.

100. This question contains Statement-1 and Statement-2. Of the four choices given after the statements, choose the one that best describes the two statements.

Statement-1: Energy is released when heavy nuclei undergo fission or light nuclei under fusion.

and

Statement-2: For heavy nuclei, binding energy per nucleon increases with increasing Z while for light nuclei it decreases with increasing Z .


- (1) Statement-1 is true, Statement-2 is true; Statement-2 is a correct explanation for Statement-1
 (2) Statement-1 is true, Statement-2 is true; Statement-2 is not a correct explanation for Statement-1
 (3) Statement-1 is true, Statement-2 is false
 (4) Statement-1 is false, Statement-2 is true

Sol: Ans [3]

Directions : Question No. 101 and 102 are based on the following paragraph.

Consider a block of conducting material of resistivity ρ shown in the figure. Current I enters at A and leaves from D . We apply superposition principle to find voltage ΔV developed between B and C . The calculation is done in the following steps :

- (i) Take current I entering from A and assume it to spread over a hemispherical surface in the block.
 (ii) Calculate field $E(r)$ at distance r from A by using Ohm's law $E = \rho J$, where J is the current per unit area at r .
 (iii) From the r dependence of $E(r)$, obtain the potential $V(r)$ at r
 (iv) Repeat (i), (ii) and (iii) for current I leaving D and superpose results for A and D .



101. For current entering at A , the electric field at a distance r from A is

- (1) $\frac{\rho I}{r^2}$ (2) $\frac{\rho I}{2\pi r^2}$ (3) $\frac{\rho I}{4\pi r^2}$ (4) $\frac{\rho I}{8\pi r^2}$

Sol: Ans [2] $J = \frac{I}{2\pi r^2}$

$$E = \frac{\rho I}{2\pi r^2}$$

102. ΔV measured between B and C is

- (1) $\frac{\rho I}{a} - \frac{\rho I}{(a+b)}$ (2) $\frac{\rho I}{2\pi a} - \frac{\rho I}{2\pi(a+b)}$ (3) $\frac{\rho I}{2\pi(a-b)}$ (4) $\frac{\rho I}{\pi a} - \frac{\rho I}{\pi(a+b)}$

Sol: Ans [4] For entering current, $v_B = \frac{\rho I \times a}{2\pi a^2} = \frac{\rho I}{2\pi a}$

$$v_C = \frac{\rho I}{2\pi(a+b)}$$

For leaving current, $v_B' = -\frac{\rho I}{2\pi(a+b)}$


$$v_C' = -\frac{\rho I}{2\pi a}$$

⇒ Potential difference between B and C

$$= \left(\frac{\rho I}{2\pi a} - \frac{\rho I}{2\pi(a+b)} \right) - \left(\frac{\rho I}{2\pi(a+b)} - \frac{\rho I}{2\pi a} \right) = \frac{\rho I}{\pi a} - \frac{\rho I}{\pi(a+b)}$$

Directions : Questions No. 103, 104 and 105 are based on the following paragraph

Wave property of electrons implies that they will show diffraction effects. Davisson and Germer demonstrated this by diffracting electrons from crystals. The law governing the diffraction from a crystal is obtained by requiring that electron waves reflected from the planes of atoms in a crystal interfere constructively (see figure)



103. If a strong diffraction peak is observed when electrons are incident at an angle i from the normal to the crystal planes with distance d between them (see figure), de Broglie wavelength λ_{dB} of electrons can be calculated by the relationship (n is an integer)

(1) $2d \cos i = n \lambda_{dB}$ (2) $2d \sin i = n \lambda_{dB}$ (3) $d \cos i = n \lambda_{dB}$ (4) $d \sin i = n \lambda_{dB}$

Sol: Ans [1] $AB = d \cos i$

Path difference = $2d \cos i$

For constructive interference, $2d \cos i = n\lambda_{dB}$

104. Electrons accelerated by potential V are diffracted from a crystal. If $d = 1 \text{ \AA}$ and $i = 30^\circ$, V should be about ($h = 6.6 \times 10^{-34} \text{ Js}$, $m_e = 9.1 \times 10^{-31} \text{ kg}$, $e = 1.6 \times 10^{-19} \text{ C}$)


(1) 50 V (2) 500 V (3) 1000 V (4) 2000 V

Sol: Ans [1] $\lambda_{dB} = \frac{h}{\sqrt{2m_e eV}}$, using (i),





$$2 \times 10^{-10} \times \cos 30 = \frac{6.6 \times 10^{-34}}{\sqrt{2 \times 9.1 \times 10^{-31} \times 1.6 \times 10^{-19} \times V}}$$

Solving, $V = 50 \text{ V}$

105. In an experiment, electrons are made to pass through a narrow slit of width d comparable to their de Broglie wavelength. They are detected on a screen at a distance D from the slit (see figure)



Which of the following graphs can be expected to represent the number of electrons N detected as a function of the detector position y ($y = 0$ corresponds to the middle of the slit) ?



Sol: Ans [3] Width of central maxima = $\left(\frac{2\lambda D}{d}\right)$

$\therefore \lambda \approx d$

\Rightarrow width of central maxima will be more than d .



READ THE FOLLOWING INSTRUCTION CAREFULLY :

1. The candidates should fill in the required particulars on the Test Booklet and Answer Sheet (Side-1)
- 2.. For writing/marking particulars on Side-2 of the Answer Sheet, use Blue/Black Ball Point Pen only.
3. The candidates should not write their Roll numbers anywhere else (except in the specified space) on the Test Booklet/Answer sheet.
4. For each **incorrect response**, **one-third** of the total marks allotted to the question would be deducted from the total score. **No deduction** from the total score, however, will be made if no response is indicated for an item in the Answer Sheet.
5. Handle the Test Booklet and Answer sheet with care, as under no circumstances (except for discrepancy in Test Booklet code and Answer Sheet Code), will another set be provided.
6. The candidates are not allowed to do any rough work or writing work on the Answer Sheet. All calculations/writing work are to be done in the space provided for this purpose in the Test Booklet itself, marked "Space for Rough Work". This space is given at the bottom of each page and in 4 pages at the end of the booklet.
7. On completion of the test, the candidates must hand over the Answer sheet to the Invigilator on duty in the Room/Hall. **However, the candidates are allowed to take away this Test Booklet with them.**
8. Each candidate must show on demand his/her Admit Card to the Invigilator.
9. No candidate, without special permission of the Superintendent or Invigilators, should leave his/her seat.
10. The candidates should not leave the Examination Hall without handing over their Answer Sheet to the Invigilator on duty and sign the Attendance Sheet again. Cases where a candidate has not signed the Attendance Sheet a second time will be deemed not to have handed over the Answer Sheet and dealt with as an unfair means case. **The candidates are also required to put their left hand THUMB impression in the space provided in the Attendance Sheet.**
11. Use of Electronic/Manual Calculate and any electronic item like mobile phone, page etc. is prohibited.
12. The candidates are governed by all Rules and Regulations of the Board with regard to their conduct in the Examination Hall. All cases of unfair means will be dealt with as per Rules and Regulations of the Board.
13. No part of the Test Booklet and Answer Sheet shall be detached under any circumstances.
14. Candidates are not allowed to carry any textual material, printed or written, bits of papers, pager, mobile phone, electronic device or any other material except the Admit Card inside the examination hall/room.

